

Cytokine and CAM Antagonists: Janus Associated Kinase (JAK) Inhibitors

Medical policy no. 66.27.00.AH-4

Effective Date: TBD

Related medical policies:

Policy Number	Policy Name
66.27.00.AA	Cytokine and CAM Antagonists: Tumor Necrosis Factor (TNF) Inhibitors
66.27.00.AB	Cytokine and CAM Antagonists: IL-4/IL-13 Inhibitors
66.27.00.AC	Cytokine and CAM Antagonists: IL-6 Inhibitors
66.27.00.AD	Cytokine and CAM Antagonists: IL-12/IL-23 Inhibitors
66.27.00.AE	Cytokine and CAM Antagonists: IL-17 Inhibitors
66.27.00.AF	Cytokine and CAM Antagonists: Oral PDE-4 Inhibitors
66.27.00.AG	Cytokine and CAM Antagonists: T-Lymphocyte Inhibitors
66.27.00.AI	Cytokine and CAM Antagonists: IL-1 Inhibitors
66.27.00.AJ	Cytokine and CAM Antagonists: Integrin Receptor Antagonists
66.27.00.AK	Cytokine and CAM Antagonists: S1-P Receptor Modulator

Note: New-to-market drugs included in this class based on the Apple Health Preferred Drug List are non-preferred and subject to this prior authorization (PA) criteria. Non-preferred agents in this class require an inadequate response or documented intolerance due to severe adverse reaction or contraindication to at least TWO preferred agents. If there is only one preferred agent in the class documentation of inadequate response to ONE preferred agent is needed. If a drug within this policy receives a new indication approved by the Food and Drug Administration (FDA), medical necessity for the new indication will be determined on a case-by-case basis following FDA labeling.

To see the list of the current Apple Health Preferred Drug List (AHPDL), please visit: <https://www.hca.wa.gov/assets/billers-and-providers/apple-health-preferred-drug-list.xlsx>

Medical necessity

Drug	Medical Necessity
abrocitinib (Cibinqo) baricitinib (Olumiant) deucravacitinib (Sotyktu) ritlecitinib (Litfulo) ruxolitinib (Opzelura) tofacitinib citrate (Xeljanz/XR) upadacitinib (Rinvoq, Rinvoq LQ)	Janus Associated Kinase (JAK) Inhibitors – abrocitinib, baricitinib, deucravacitinib, ritlecitinib, ruxolitinib, tofacitinib, upadacitinib may be considered medically necessary in patients who meet the criteria described in the clinical policy below. If all criteria are not met, the clinical reviewer may determine there is a medically necessary need and approve on a case-by-case basis. The clinical reviewer may choose to use the reauthorization criteria when a patient has been previously established on therapy and is new to Apple Health.

Clinical policy:

Clinical Criteria	
<p>Alopecia areata baricitinib (Olumiant) ritlecitinib (Litfulo)</p>	<p>baricitinib (Olumiant) or ritlecitinib (Litfulo) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. The patient meets the age appropriate limit for the requested product: <ol style="list-style-type: none"> a. For baricitinib: 18 years of age or older; OR b. For ritlecitinib: 12 years of age or older; AND 2. Prescribed by, or in consultation with, a dermatologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of severe alopecia areata; AND 5. Current episode of alopecia areata lasting more than 6 months; AND 6. Patient has ≥50% of the scalp hair loss (Severity of Alopecia Tool [SALT] score >50%); AND 7. History of failure to one of the following unless all are contraindicated or not tolerated: <ol style="list-style-type: none"> a. High-potency topical corticosteroids [minimum trial of 6 weeks]; OR b. Intralesional corticosteroids [minimum trial of 6 weeks]; OR c. Systemic therapy (i.e., oral corticosteroids, methotrexate, cyclosporine) [minimum trial of 6 weeks]. <p>If ALL criteria are met, the request will be authorized for 6 months.</p>
Criteria (Reauthorization)	
	<p>baricitinib (Olumiant) or ritlecitinib (Litfulo) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND 2. Documentation is submitted demonstrating a positive clinical response (e.g. improvement in SALT score). <p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Ankylosing Spondylitis tofacitinib (Xeljanz) upadacitinib (Rinvoq)</p> <p>Non-Radiographic Axial Spondyloarthritis Upadacitinib (Rinvoq)</p>	<p>tofacitinib (Xeljanz) or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older, AND 2. Prescribed by, or in consultation with a rheumatologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Patient meets one of the following diagnosis criteria: <ol style="list-style-type: none"> a. For tofacitinib: Diagnosis of Ankylosing Spondylitis (AS); OR

	<ul style="list-style-type: none"> b. For upadacitinib: Diagnosis of ankylosing spondylitis (AS) or non-radiographic axial spondyloarthritis; AND 5. High disease activity as indicated by a Bath Ankylosing Disease Activity Index (BASDAI) score of at least 4 or an Ankylosing Spondylitis Disease Activity Score (ASDAS) score of at least 2.1; AND 6. Treatment with at least two different NSAIDs (e.g., indomethacin, meloxicam, celecoxib, naproxen, nabumetone, etc.) have been ineffective unless all are contraindicated or not tolerated [minimum trial of four weeks] 7. Disease manifested as either of the following: <ul style="list-style-type: none"> a. Axial disease; OR b. Peripheral arthritis; AND <ul style="list-style-type: none"> i. Treatment with at least one non-Cytokine and CAM disease-modifying antirheumatic drug (DMARD) (e.g., methotrexate, sulfasalazine, leflunomide) has been ineffective unless all are contraindicated or not tolerated [minimum trial of 3 months]. 8. Treatment with two preferred Cytokine & CAM Apple Health Preferred Drug List (AHPDL) medications have each been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 weeks]. <p>If ALL criteria are met, the request will be authorized for 6 months.</p>
Criteria (Reauthorization)	
<p>tofacitinib (Xeljanz) or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ul style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., decrease in BASDAI or ASDAS score). <p>If ALL criteria are met, the request will be authorized for 12 months.</p>	
<p>Atopic Dermatitis abrocitinib (Cibinqo) ruxolitinib (Opzelura) upadacitinib (Rinvoq)</p>	<p>abrocitinib (Cibinqo), ruxolitinib (Opzelura), or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ul style="list-style-type: none"> 1. Patient is 12 years of age or older; AND 2. Prescribed by, or in consultation with an allergist, dermatologist, or an immunologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of one of the following: <ul style="list-style-type: none"> a. For abrocitinib and upadacitinib, moderate to severe atopic dermatitis; OR b. For ruxolitinib, mild to moderate atopic dermatitis; AND 5. For upadacitinib, the patient weighs 40 kg or more; AND

6. Documentation is provided demonstrating:
 - a. For abrocitinib and upadacitinib, one of the following:
 - i. Body surface area (BSA) involvement of at least 10% unless there is involvement of sensitive skin areas such as hands, feet, face, neck, genitalia, or intertriginous areas; **OR**
 - ii. Baseline disease severity scale scoring supporting diagnosis of moderate to severe atopic dermatitis (e.g., Investigator’s Global Assessment (IGA) score of 3 or greater; Eczema Area and Severity Index (EASI), Patient Oriented Eczema Measure (POEM), etc.); **OR**
 - b. For ruxolitinib:
 - i. BSA involvement does not exceed 20%; **AND**
7. Patient is experiencing functional impairment due to atopic dermatitis, which may include, but is not limited to:
 - a. Activities of daily living (ADLs); **OR**
 - b. Skin infections; **OR**
 - c. Sleep disturbances; **AND**
8. History of failure, defined as the inability to achieve or maintain remission to at **LEAST TWO** of the following groups unless all are contraindicated or not tolerated [minimum trial of 28-days each]:
 - a. Group 1: Topical corticosteroids of at least medium/moderate potency (e.g. betamethasone, clobetasol, halobetasol, hydrocortisone, mometasone)
 - b. Group 2: Topical calcineurin inhibitors (e.g. pimecrolimus cream, tacrolimus ointment)
 - c. Group 3: Topical PDE-4 inhibitors (e.g. crisaborole); **AND**
9. For abrocitinib and upacitinib, treatment with dupilumab (Dupixent) has been ineffective, contraindicated, or not tolerated [minimum trial of 16 weeks].

If ALL criteria are met, the request will be authorized for **6 months**.

Criteria (Reauthorization)

abrocitinib (Cibinqo), upadacitinib (Rinvoq), or ruxolitinib (Opzelura) may be approved when all the following documented criteria are met:

1. Not used in combination with another Cytokine and CAM medication; **AND**
2. Documentation is submitted demonstrating disease stability, or a positive clinical response defined by BOTH of the following:
 - a. At least ONE of the following:
 - i. Reduction in body surface area involvement of at least 20% compared to baseline; **OR**
 - ii. Achieved or maintained clear or minimal disease from baseline (equivalent to IGA score of 0 or 1); **OR**

	<p>iii. Experienced or maintained a decrease in EASI score of at least 50% compared to baseline; AND</p> <p>b. An improvement in functional impairment (e.g., improvement in ADLs, skin infections, or sleep disturbance).</p> <p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Crohn's Disease upadacitinib (Rinvoq)</p>	<p>upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older, AND 2. Prescribed by, or in consultation with a gastroenterologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of moderate to severe Crohn's disease (CD); AND <ol style="list-style-type: none"> a. Treatment with conventional therapy has been ineffective unless all are contraindicated or not tolerated. Conventional therapy is defined as: <ol style="list-style-type: none"> i. Oral corticosteroids (e.g., prednisone, methylprednisolone) used short-term to induce remission or alleviate signs/symptoms of disease flare; AND ii. At least one immunomodulatory agent (e.g., methotrexate, azathioprine, 6-mercaptopurine) [minimum trial of 12 weeks]; OR b. Documentation of high-risk disease (e.g., symptoms despite conventional therapy, obstruction, abscess, stricture, phlegmon, fistulas, resection, extensive bowel involvement, early age of onset, growth retardation, Crohn's Disease Activity Index (CDAI) > 450, Harvey-Bradshaw index > 7); AND 5. Treatment with adalimumab has been ineffective, contraindicated, or not tolerated [minimum trial of 12 weeks]. <p>If ALL criteria are met, the request will be authorized for 6 months.</p> <p>Criteria (Reauthorization)</p> <p>upadacitinib (Rinvoq) may be approved when all of the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., improvement in endoscopic activity, taper or discontinuation of corticosteroids, reduction in number of liquid stools, decrease in presence and severity of abdominal pain, decrease in CDAI, decrease in Harvey-Bradshaw index).

	<p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Plaque Psoriasis deucravacitinib (Sotyktu)</p>	<p>deucravacitinib (Sotyktu) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older, AND 2. Prescribed by, or in consultation with a dermatologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of moderate to severe plaque psoriasis; AND 5. Presence of ongoing disease for greater than 6 months; AND 6. The patient meets one of the following: <ol style="list-style-type: none"> a. Disease affects at least 10% body surface area; OR b. Disease affects the face, ears, hands, feet, or genitalia; AND 7. Baseline assessments are included (e.g., body surface area (BSA), Psoriasis Are and Severity Index (PASI), Psoriasis Physician’s Global Assessment (PGA), itch numeric rating scale, etc.); AND 8. History of failure to one of the following unless all are contraindicated or not tolerated: <ol style="list-style-type: none"> a. Phototherapy (UVB or PUVA) [minimum trial of 12 weeks]; OR b. Treatment with at least one non-Cytokine and CAM DMARD (e.g., methotrexate, cyclosporine, acitretin, azathioprine, etc.) [minimum trial of 12 weeks]; AND 9. Treatment with two preferred Cytokine & CAM Apple Health Preferred Drug List (AHPDL) medications have each been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 weeks]. <p>If ALL criteria are met, the request will be authorized for 6 months.</p> <p>Criteria (Reauthorization)</p> <p>deucravacitinib (Sotyktu) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., improvement in BSA, PASI, Psoriasis PGA, itch numeric rating scale). <p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Polyarticular Juvenile Idiopathic Arthritis tofacitinib (Xeljanz) Upadacitinib (Rinvoq, Rinvoq LQ)</p>	<p>Tofacitinib (Xeljanz) or upadacitinib (Rinvoq, Rinvoq LQ) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 2 to 17 years of age or older, AND 2. Prescribed by, or in consultation with a rheumatologist; AND

	<ol style="list-style-type: none"> 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of Polyarticular Juvenile Idiopathic Arthritis (PJIA); AND 5. Documentation of current weight is provided; AND 6. Treatment with at least one non-Cytokine and CAM DMARD (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine, azathioprine, cyclosporine) has been ineffective unless all are contraindicated or not tolerated [minimum trial of 3 months]; AND 7. Treatment with two preferred Cytokine & CAM Apple Health Preferred Drug List (AHPDL) medications have each been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 weeks]. <p>If ALL criteria are met, the request will be authorized for 6 months.</p> <p>Criteria (Reauthorization)</p> <p>Tofacitinib (Xeljanz) or upadacitinib (Rinvoq, Rinvoq LQ) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., improvement in joint pain, swelling, activities of daily living, reduction in diseases flares, etc.). <p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Psoriatic Arthritis tofacitinib (Xeljanz) upadacitinib (Rinvoq, Rinvoq LQ)</p>	<p>tofacitinib (Xeljanz) or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. The patient meets the appropriate age limit for the requested product: <ol style="list-style-type: none"> a. For tofacitinib: 18 years of age or older; OR b. For upadacitinib: 2 years of age or older; AND 2. Prescribed by, or in consultation with a rheumatologist or dermatologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of Psoriatic Arthritis (PsA); AND 5. For pediatric upadacitinib requests, documentation of current weight is provided; AND 6. Patient meets one of the following: <ol style="list-style-type: none"> a. Treatment with at least one non-Cytokine and CAM DMARD (e.g., methotrexate, sulfasalazine, leflunomide, cyclosporine) has been ineffective unless all are

	<p>contraindicated or not tolerated [minimum trial of 3 months]; OR</p> <p>b. Presence of active, severe disease as indicated by provider assessment and the presence of at least <u>ONE</u> of the following:</p> <ol style="list-style-type: none"> i. Erosive disease; OR ii. Elevated C-reactive protein (CRP) or erythrocyte sedimentation rate (ESR); OR iii. Long-term damage interfering with function (e.g., joint deformities, vision loss); OR iv. Major impairment of quality of life due to high disease activity at many sites (including dactylitis, enthesitis) or functionally limiting arthritis at a few sites; AND <p>7. Treatment with two preferred Cytokine & CAM Apple Health Preferred Drug List (AHPDL) medications have each been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 weeks].</p> <p>If ALL criteria are met, the request will be authorized for 6 months.</p> <p>Criteria (Reauthorization)</p> <p>Tofacitinib (Xeljanz) or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., improvement in joint pain, swelling, activities of daily living, reduction in diseases flares, etc.). <p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Rheumatoid Arthritis baricitinib (Olumiant) tofacitinib (Xeljanz) upadacitinib (Rinvoq)</p>	<p>baricitinib (Olumiant), tofacitinib (Xeljanz) or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older, AND 2. Prescribed by, or in consultation with a rheumatologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of Rheumatoid Arthritis (RA); AND 5. Baseline assessments are included (e.g., Disease Activity Score for 28 joints (DAS28) with the CRP, DAS28 with ESR, Simplified Disease Activity Index (SDAI), Clinical Disease Activity Index (CDAI), Routine Assessment of Patient Index Data 3 (RAPID3), Patient Activity Scale (PAS) II); AND 6. Treatment with at least one non-Cytokine and CAM DMARD (e.g., methotrexate, sulfasalazine, hydroxychloroquine,

	<p>leflunomide, cyclosporine, azathioprine) has been ineffective unless all are contraindicated or not tolerated [minimum trial of 3 months]; AND</p> <p>7. Treatment with two preferred Cytokine & CAM Apple Health Preferred Drug List (AHPDL) medications have each been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 weeks].</p> <p>If ALL criteria are met, the request will be authorized for 6 months.</p> <p>Criteria (Reauthorization)</p> <p>baricitinib (Olumiant), tofacitinib (Xeljanz), or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND 2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g. improvement in DAS28 with CRP/ESR, SDAI, CDAI, RAPID3, PAS II scores). <p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Ulcerative Colitis tofacitinib (Xeljanz) upadacitinib (Rinvoq)</p>	<p>tofacitinib (Xeljanz) or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 18 years of age or older, AND 2. Prescribed by, or in consultation with a gastroenterologist; AND 3. Not used in combination with another Cytokine and CAM medication; AND 4. Diagnosis of moderate-to-severe Ulcerative Colitis (UC); AND 5. Baseline assessments are included (e.g., stool frequency, endoscopy results, presence of rectal bleeding, disease activity scoring tool); AND 6. Treatment with conventional therapy (e.g., systemic corticosteroids, azathioprine, mesalamine, sulfasalazine) has been ineffective unless all are contraindicated or not tolerated [minimum trial of 12 weeks]; AND 7. Treatment with adalimumab has been ineffective, contraindicated, or not tolerated [minimum trial of 12 weeks]. <p>If ALL criteria are met, the request will be authorized for 6 months.</p> <p>Criteria (Reauthorization)</p> <p>tofacitinib (Xeljanz) or upadacitinib (Rinvoq) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication; AND

	<p>2. Documentation is submitted demonstrating disease stability or a positive clinical response (e.g., decreased stool frequency, decreased rectal bleeding, improvement in endoscopic activity, tapering or discontinuation of corticosteroid therapy, or improvement on a disease activity scoring tool).</p> <p>If ALL criteria are met, the request will be authorized for 12 months.</p>
<p>Vitiligo ruxolitinib (Opzelura)</p>	<p>ruxolitinib (Opzelura) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Patient is 12 years of age or older; AND 2. Prescribed by, or in consultation with, a dermatologist; AND 3. Not used in combination with another Cytokine and CAM medication or oral immunosuppressants (e.g. azathioprine, cyclosporine); AND 4. Diagnosis of nonsegmental vitiligo including: <ol style="list-style-type: none"> a. Patient has chronic disease (greater than 3 months); AND b. Provider attests vitiligo involves a total body surface area (BSA) that is 10% or less; AND 5. Treatment with at least one therapy from two of the following categories has been ineffective or not tolerated. If one category is contraindicated, the patient must still try one therapy from two categories: <ol style="list-style-type: none"> a. Phototherapy b. Topical calcineurin inhibitors (e.g. tacrolimus) c. Medium to high potency topical corticosteroids (e.g. betamethasone, mometasone, clobetasol, fluocinonide) <p>If ALL criteria are met, the request will be authorized for 6 months.</p> <p>Criteria (Reauthorization)</p> <p>ruxolitinib (Opzelura) may be approved when all the following documented criteria are met:</p> <ol style="list-style-type: none"> 1. Not used in combination with another Cytokine and CAM medication or oral immunosuppressant; AND 2. Documentation is submitted demonstrating a positive clinical response (e.g. improvement in F-VASI and/or T-VASI score from baseline, or reduction in total BSA involvement) <p>If ALL criteria are met, the request will be authorized for 12 months.</p>

Dosage and quantity limits

Drug	Indication	FDA Approved Dosing	Dosage Form and Quantity Limit
Cibinqo	Atopic Dermatitis	100-200 mg once daily	<ul style="list-style-type: none"> • 50 mg tablet: 30 tablets per 30 days • 100 mg tablet: 30 tablets per 30 days • 200 mg tablet: 30 tablets per 30 days
Litfulo	Alopecia Areata	50 mg once daily	<ul style="list-style-type: none"> • 50 mg capsules: 30 capsules per 30 days
Olumiant	Alopecia Areata	2-4 mg once daily	

	Rheumatoid Arthritis	2 mg once daily	<ul style="list-style-type: none"> 1 mg tablet: 30 tablets per 30 days 2 mg tablet: 30 tablets per 30 days 4 mg tablet: 30 tablets per 30 days
Opzelura	Atopic Dermatitis	Apply a thin layer topically twice daily	<ul style="list-style-type: none"> Max 60 grams per 1 week OR 100 grams per 2 weeks
	Vitiligo	Apply a thin layer topically twice daily	
Rinvoq	Ankylosing Spondylitis	15 mg once daily	<ul style="list-style-type: none"> 1 mg/mL solution: 240 mL per 30 days* 15 mg tablet: 30 tablets per 30 days 30 mg tablet: 30 tablets per 30 days 45 mg tablet: 30 tablets per 30 days
	Atopic Dermatitis	Less than 65 years of age: 15-30 mg once daily 65 years of age and older: 15 mg once daily	
	Crohn's Disease	Induction: 45 mg once daily for 12 weeks Maintenance: 15-30 mg once daily	
	Polyarticular juvenile idiopathic arthritis	10-20kg: 3 mg (oral solution) twice daily 20 to less than 30 kg: 4 mg (oral solution) twice daily 30 kg or greater: 6 mg twice daily (oral solution) or 15 mg (tablet) once daily	
	Psoriatic Arthritis	10-20kg: 3 mg (oral solution) twice daily 20 to less than 30 kg: 4 mg (oral solution) twice daily 30 kg or greater: 6 mg twice daily (oral solution) or 15 mg once daily	
	Non-radiographic axial spondyloarthritis	15 mg once daily	
	Rheumatoid Arthritis	15 mg once daily	
	Ulcerative Colitis	Induction: 45 mg once daily for 8 weeks Maintenance: 15-30 mg once daily	
Sotyktu	Plaque psoriasis	6 mg once daily	<ul style="list-style-type: none"> 6 mg tablet: 30 tablets per 30 days
Xeljanz, Xeljanz XR	Ankylosing Spondylitis	5 mg twice daily 11 mg once daily	<ul style="list-style-type: none"> 1 mg/mL solution: 240 mL per 30 days* 5 mg tablet: 60 tablets per 30 days 10 mg tablet: 60 tablets per 30 days 11 mg XR tablet: 30 tablets per 30 days 22 mg XR tablet: 30 tablets per 30 days
	Polyarticular Juvenile Idiopathic Arthritis	10 to less than 20 kg: 3.2 mg twice daily 20kg to less than 40 kg: 4 mg twice daily 40kg or more: 5 mg twice daily	
	Psoriatic Arthritis	5 mg twice daily 11 mg once daily	
	Rheumatoid Arthritis	5 mg twice daily 11 mg once daily	

	Ulcerative Colitis	<u>IR Tablet</u> Induction: 10 mg twice daily for up to 16 weeks Maintenance: 5 mg twice daily <u>ER Tablet</u> Induction: 22 mg once daily for up to 16 weeks Maintenance: 11 mg once daily	
--	--------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--

*Dosing for PJI is based on body weight. Patients on Rinvoq with body weight greater than >30 kg may be switched to Rinvoq 15 mg tablets. Patients on Xeljanz with body weight >40kg may be switched to Xeljanz 5 mg tablets.

Coding:

HCPCS Code	Description
N/A	N/A

Background:

Alopecia Areata

There are no formal U.S. treatment guidelines addressing alopecia areata. The [2020 Alopecia Areata Consensus of Experts \(ACE\)](#) study provides international expert consensus statements aiming to help medical practitioners select optimal alopecia areata management strategies. ACE consensus for topical treatments in alopecia areata state topical corticosteroids can be prescribed as first-line topical treatment (alone or in combination) to treat scalp, eyebrow, or beard alopecia areata. ACE notes that in adults, the most appropriate first-line treatment, when Severity of Alopecia Tool (SALT) score is greater than 50%, is topical or oral corticosteroids. Steroid-sparing agents are commonly used to mitigate the risk of adverse effects associated with prolonged use of high-dose systemic corticosteroids. In adults with alopecia areata, cyclosporine is an effective monotherapy agent. It was acknowledged that methotrexate is sometimes used as monotherapy in severe alopecia areata. Consensus was not achieved in any questions regarding use of azathioprine. Treatment with a JAK inhibitor is noted as a preferred second line agent.

Ankylosing spondylitis and non-radiographic axial spondyloarthritis

The [2019 American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network \(ACR/SAA/SPARTAN\)](#) guidelines on the treatment of ankylosing spondylitis strongly recommend the use of NSAIDs as first-line treatment (with 70-80% responding). Recommendations against the use of non-biologic DMARDs are made for patients with active ankylosing spondylitis despite NSAID treatment. Some benefit has been seen in patients with peripheral arthritis, thus treatment with sulfasalazine or methotrexate may be considered in patients with predominantly peripheral disease; however, evidence is based on older RCTs with very low quality of evidence. For those patients with inadequate response despite continuous NSAID treatment, the ACR strongly recommends use of TNF inhibitors over no treatment with TNF inhibitors. In patients with secondary nonresponse to TNF inhibitors, the guidelines conditionally recommend treatment with a different TNF inhibitor over treatment with a non-TNF inhibitor biologic. The [2022 Assessment of SpondyloArthritis international Society \(ASAS\)-EULAR](#) guidelines for the treatment of axial spondyloarthritis (axSpA) reference the use of JAK inhibitors in the treatment algorithm. The term axial spondyloarthritis (axSpA), encompasses both active ankylosing spondylitis (or radiographic AS) and nr-axSpA as one entity part of the same chronic inflammatory musculoskeletal spectrum with similar clinical presentations, comorbidities, disease burden, and treatment response. ASAS/EULAR recommends patients try and fail at least 2 NSAIDs over 4 weeks as first line therapy and treat local musculoskeletal inflammation with glucocorticoid injection; sulfasalazine may be considered in patients with peripheral symptoms, however use of conventional non-biologic DMARDs (e.g. sulfasalazine, leflunomide, methotrexate, etc.) is not recommended in axial disease. In contrast to ACR/SAA/SPARTAN, ASAS/EULAR guidelines highly recommend treatment with a TNF inhibitor, IL-17 inhibitor, or JAK inhibitor for patients with high disease activity, defined by a BASDAI of at least 4 or an ASDAS of at least 2.1, despite conventional treatment with NSAIDs. Starting with a TNF inhibitor or IL-17 inhibitor is preferred clinically, given long term data for use of JAK inhibitors in axSpA is still missing. There is no specific treatment algorithm after primary non-response to biologic (TNF inhibitor or IL-17 inhibitor) or JAK inhibitor therapy.

Atopic dermatitis

Treatments for mild-to-moderate AD include topical corticosteroids (TCS), topical calcineurin inhibitors (TCI), phototherapy, and crisaborole (Eucrisa), and Opzelura (ruxolitinib). Symptomatic treatments include oral and topical antihistamines and sleep aids for nighttime pruritus. Treatment choice between these products is dependent on severity, location, and other patient specific factors (e.g., allergies, age). According to [American Academy of Dermatology](#) (AAD) guidelines, TCIs may be preferable to TCS in patients with recalcitrance to steroids, sensitive areas involved, steroid-induced atrophy, and long-term uninterrupted topical steroid use. Treatment for moderate to severe disease not amenable to topicals includes systemic immunosuppressants (e.g., corticosteroids, cyclosporine, methotrexate, azathioprine, mycophenolate mofetil), JAK inhibitors (e.g., abrocitinib, upadacitinib), and dupilumab (Dupixent). Currently, there are no head- to-head trials evaluating safety and/or efficacy differences or superiority between biologic therapies in atopic dermatitis.

Crohn's Disease

Therapeutic recommendations for patients with Crohn's disease (CD) are established based upon disease location, disease severity, disease associated complications, and future disease prognosis. The goals of therapy are to induce remission, prevent relapse, and prevent occurrence of disease complications, such as stricture and fistula. According to the [2018 American College of Gastroenterology \(ACG\)](#) guidelines, for patients with moderate to severe disease and those with moderate to high-risk disease treatment with oral corticosteroids used short term to induce remission is recommended (strong recommendation, moderate level of evidence). However, it is noted that one in five patients will become steroid refractory which is thought to be the result of unreliable efficacy in healing of the mucosa associated with steroids (weak recommendation, low level of evidence). Corticosteroids are also implicated in the development of perforating complications (abscess and fistula) and are relatively contraindicated in those patients. The [2021 American Gastroenterological Association \(AGA\)](#) clinical guidelines make similar recommendations and suggest the use of corticosteroids in adult outpatients with moderate to severe CD over no treatment for induction of remission (conditional recommendation, moderate level of evidence). In patients with moderate to severe CD who remain symptomatic despite current or prior corticosteroid therapy, 2018 ACG guidelines recommend immunomodulators such as azathioprine, 6-mercaptopurine (strong recommendation, moderate level of evidence), and methotrexate (conditional recommendation, low level of evidence) to be effective for maintenance of remission. Due to slow time to clinical response that may not be evident for as long as 12 weeks, these agents are not recommended for short-term induction. The 2021 AGA guidelines make similar suggestions and recommend use of thiopurines over no treatment for the maintenance of remission (conditional recommendation, low level of evidence). The timing of introduction of biologic agents is a matter of debate and more studies are needed to assess stepwise approach versus earlier administration of biologic agents in patients with moderate to severe disease. The [2019 British Society of Gastroenterology](#) guidelines suggest that systemic corticosteroids are still an effective initial therapy for uncomplicated luminal moderate to severe disease, regardless of disease location; however, every effort should be made to limit exposure (strong recommendation, high-quality evidence). In patients with an aggressive disease course, or high risk, poor prognostic factors, early introduction of biologics may be considered (weak recommendation, moderate-quality evidence). High risk features include extensive disease, complex (stricturing or penetrating disease), perianal fistulizing disease, age under 40 years at diagnosis, and the need for steroids to control index flare; however, the predictive power of these features is limited.

Plaque Psoriasis

Joint American Academy of Dermatology–National Psoriasis Foundation guidelines for the [management of psoriasis with systemic nonbiologic therapies](#) and for the [management and treatment of psoriasis with biologics](#) indicate that the majority of patients are capable of adequately controlling disease solely with topical medications or phototherapy. Phototherapy is recognized as a beneficial therapy for controlled plaque psoriasis and is a cost-effective treatment strategy. Additionally, oral immunomodulatory medications (e.g., methotrexate, cyclosporine, acitretin) are cost-effective therapies with a well-known safety profile for the treatment of plaque psoriasis. For moderate-to-severe disease, where a JAK inhibitor or biologics are warranted, deucravacitinib (Sotyktu) is one of many options. However, it would not be indicated for mild psoriasis given that patients are better managed from a safety perspective on well-established therapies (e.g., topical agents, phototherapy, conventional DMARDS, apremilast [Otezla]).

Polyarticular Juvenile Idiopathic Arthritis (PJIA)

Juvenile idiopathic arthritis (JIA) is a grouping of inflammatory disorders that affect children. Polyarticular juvenile idiopathic arthritis (PJIA) is a subset of JIA, which is defined by the presence arthritis in five or more joints during the first six months of illness. Other subsets of JIA include ERA, oligoarthritis (less than five joints affected), systemic juvenile idiopathic arthritis (SJIA; fever, rash, hepatic/splenic/lymphatic involvement) and psoriatic arthritis (psoriasis and dactylitis). While these are distinct disease states, their pathogenesis and presentation are similar so there is significant overlap in effective treatments. The [2019 American College of Rheumatology/Arthritis Foundation](#) (ACR) guidelines for non-systemic polyarthritis (PJIA) strongly recommend initial therapy with a DMARD for all patients with JIA and active polyarthritis; methotrexate has the strongest evidence, but sulfasalazine and leflunomide can also be used. Regardless of disease activity, initial therapy with a DMARD is recommended over a biologic, though there may be certain situations where a biologic as initial therapy is preferred (i.e., high risk joints such as cervical spine, wrist, or hip involved). For patients with continued moderate to high disease activity, the guidelines recommend adding a TNF inhibitor, abatacept, or tocilizumab as second-line.

Psoriatic Arthritis

The [2018 American College of Rheumatology/National Psoriasis Foundation](#) (ACR) guidelines for psoriatic arthritis make a conditional recommendation for starting a TNF inhibitor over an oral small molecule (OSM) as a first-line option for patients who are treatment-naïve with active psoriatic arthritis. This recommendation is based on low- to very-low quality of evidence. Many of the studies in which greater benefit was seen in terms of disease severity or radiographic progression compared methotrexate to TNF inhibitors, however, most patients included in these groups were not truly treatment naïve to OSM medications. Guidelines note that OSM can be used first-line in naïve patients who do not have severe PsA, severe PsO, prefers oral therapy, or has contraindications to TNF inhibitors. In patients who continue to have active disease despite OSM treatment, it is recommended to switch to a TNF inhibitor rather than trying a different OSM. The 2018 ACR guidelines for psoriatic arthritis also conditionally recommend for use of a TNF inhibitor biologics over IL-17 inhibitors (ixekizumab, secukinumab) or IL-12/23 inhibitors (ustekinumab).

Rheumatoid Arthritis

The [2021 American College of Rheumatology \(ACR\)](#) guidelines for rheumatoid arthritis strongly recommend the use of conventional synthetic disease-modifying antirheumatic drug (csDMARD) monotherapy (methotrexate preferred) in patients who are DMARD-naïve with moderate-to-severe RA. Recommended csDMARDs include methotrexate, sulfasalazine, hydroxychloroquine, and leflunomide. Despite moderate evidence in the SELECT-EARLY study noting higher efficacy of upadacitinib over methotrexate in DMARD-naïve patients with moderate-to-severe RA, there is limited long-term safety data to strongly recommend the use of tsDMARDs (e.g., JAK inhibitors) as first line therapy. Therefore, methotrexate monotherapy remains the preferred first-line therapy over tsDMARDs in DMARD-naïve patients based on established safety and efficacy. Additionally, JAK inhibitors are not FDA approved for use in csDMARD-naïve patients. The [2019 European League Against Rheumatism \(EULAR\)](#) guidelines follow similar recommendations to the 2021 ACR guidelines, and state that patients with highly active RA despite treatment with csDMARDs may receive a biologic DMARD or JAK inhibitor based on high level of evidence.

Ulcerative Colitis

The [2019 American College of Gastroenterology \(ACG\)](#) clinical guideline on the management of ulcerative colitis in adults recommend oral systemic corticosteroids for induction of remission in moderate to severe disease (strong recommendation, moderate quality of evidence). TNF inhibitors (adalimumab, golimumab, and infliximab), vedolizumab (Entyvio), and tofacitinib (Xeljanz) are also recommended for induction of remission (strong recommendation, moderate quality of evidence). For maintenance of remission, thiopurines are recommended if remission was achieved after corticosteroid induction (conditional recommendation, low quality of evidence). The guidelines note a systematic review of 1,632 patients with ulcerative colitis demonstrated that azathioprine and mercaptopurine had a 76% mean efficacy in maintaining remission. If remission was achieved with anti-TNF therapy, vedolizumab (Entyvio), or tofacitinib (Xeljanz), clinical guidelines support continuing with the same agent to maintain remission (strong recommendation, moderate quality of evidence). The [2020 American Gastroenterology Association \(AGA\)](#) guidelines make similar recommendations. Additionally, AGA recommends early use of biologic agents, rather than gradual step up after failure of 5-ASA in moderate to severe disease at high risk for colectomy. However, overall quality of evidence supporting this recommendation was rated as very low. Guidelines also note that for patients with less severe disease, 5-ASA therapy may still be a reasonable choice of therapy to start with. For maintenance of remission, AGA makes no recommendation in favor of, or against, using biologic monotherapy, rather than thiopurine monotherapy due to absence of evidence.

Vitiligo

The [2023 Position Statement from the International Vitiligo Task Force](#) recommend topical corticosteroids, topical calcinurin inhibitors (TCIs), phototherapy, and ruxolitinib as treatment options for stable vitiligo. TCIs used in combination with UV phototherapy appear to be more effective than treatment with a single agent. Ruxolitinib (Opzelura) is the first topical JAK inhibitor approved for the treatment of nonsegmental vitiligo. It may be used for vitiligo affecting up to 10% body surface area. In phase three clinical trials, ruxolitinib over 30% of participants achieved a 75% reduction in facial vitiligo area scoring index (F-VASI) compared to less than 10% with the placebo.

References

1. Cibinqo [Prescribing Information]. New York, NY; Pfizer Inc., Updated February 2023.
2. Olumiant [Prescribing Information]. Indianapolis, IN; Eli Lilly. Updated June 2022.
3. Sotyktu [Prescribing Information]. Princeton, NJ; Bristol-Myers Squibb. Updated September 2022.
4. Xeljanz/Xeljanz XR [Prescribing Information]. New York, NY; Pfizer Inc., Updated January 2022.
5. Rinvoq [Prescribing Information]. North Chicago, IL; AbbVie. Updated May 2023.
6. Fraenkel L, Bathon JM, England BR, et al. 2021 American college of rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Care Res.* 2021;73(7):924-939.
7. Opzelura [Prescribing Information]. Wilmington, DE; Incyte Corp. Updated August, 2024
8. Alten R, Mischkewitz M. 2021 ACR guideline reflects changes in RA treatment. *Nat Rev Rheumatol.* 2021;17(9):513-514. doi:10.1038/s41584-021-00667-2
9. Van Vollenhoven, R. et al. Efficacy and safety of upadacitinib monotherapy in methotrexate-naive patients with moderately-to-severely active rheumatoid arthritis (SELECT-EARLY): a multicenter, multi-country, randomized, double-blind, active-comparator-controlled trial. *Arthritis Rheumatol.* 72, 1607–1620 (2020)
10. UpToDate, Inc. General principles and overview of management of rheumatoid arthritis in adults . UpToDate [database online]. Waltham, MA. Last updated October 18, 2021. Available at: <http://www.uptodate.com/home/index.html>.
11. Smolen JS, Landewé RBM, Bijlsma JWJ, et al. EULAR recommendations for the management of rheumatoid arthritis with synthetic and biological disease-modifying antirheumatic drugs: 2019 update. *Ann Rheum Dis.* 2020;79:685-699.

12. UpToDate, Inc. Spondyloarthritis in children. UpToDate [database online]. Waltham, MA. Last updated December 4, 2020. Available at uptodate.com. Accessed February 4, 2022.
13. Ringold S, Angeles-han ST, Beukelman T, et al. 2019 American College of Rheumatology/Arthritis Foundation Guideline for the Treatment of Juvenile Idiopathic Arthritis: Therapeutic Approaches for Non-Systemic Polyarthritis, Sacroiliitis, and Enthesitis. *Arthritis Care Res (Hoboken)*. 2019.
14. UpToDate, Inc. Polyarticular juvenile idiopathic arthritis: treatment. UpToDate [database online]. Waltham, MA. Last updated October 19, 2020. Available at: <http://www.uptodate.com/home/index.html>. Last accessed November 22, 2021.
15. Singh JA, Guyatt G, Ogdie A, et al. Special Article: 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis Rheumatol*. 2019;71(1):5-32.
16. Kingsley GH, Scott DL. Assessing the effectiveness of synthetic and biologic disease-modifying antirheumatic drugs in psoriatic arthritis - a systematic review. *Psoriasis (Auckl)*. 2015;5:71-81.
17. UpToDate, Inc. Treatment of psoriatic arthritis. UpToDate [database online]. Waltham, MA. Last updated November 20, 2020. Available at: <http://www.uptodate.com/home/index.html>.
18. Ward, M.M., Deodhar, A., Gensler, L.S, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Rheumatol*, 71: 1599-1613.
19. Ramiro S, Nikiphorou E, Sepriano A, et al. ASAS-EULAR recommendations for the management of axial spondyloarthritis: 2022 update. *Ann Rheum Dis*. Published online October 21, 2022:ard-2022-223296.
20. UpToDate, Inc. Clinical manifestations of axial spondyloarthritis (ankylosing spondylitis and nonradiographic axial spondyloarthritis) in adults. UpToDate [database online]. Waltham, MA. Last updated November 2, 2022. Available at: <http://www.uptodate.com/home/index.html>.
21. UpToDate, Inc. Treatment of axial spondyloarthritis (ankylosing spondylitis and nonradiographic axial spondyloarthritis) in adults. UpToDate [database online]. Waltham, MA. Last updated August 24, 2022. Available at: <http://www.uptodate.com/home/index.html>.
22. UpToDate, Inc. Treatment of peripheral spondyloarthritis. UpToDate [database online]. Waltham, MA. Last updated March 17, 2022. Available at: <http://www.uptodate.com/home/index.html>.
23. Ward, M.M., Deodhar, A., Gensler, L.S, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the treatment of ankylosing spondylitis and nonradiographic axial spondyloarthritis. *Arthritis Rheumatol*, 71: 1599-1613.
24. Ramiro S, Nikiphorou E, Sepriano A, et al. ASAS-EULAR recommendations for the management of axial spondyloarthritis: 2022 update. *Ann Rheum Dis*. Published online October 21, 2022:ard-2022-223296.
25. UpToDate, Inc. Clinical manifestations of axial spondyloarthritis (ankylosing spondylitis and nonradiographic axial spondyloarthritis) in adults. UpToDate [database online]. Waltham, MA. Last updated November 2, 2022. Available at: <http://www.uptodate.com/home/index.html>.
26. UpToDate, Inc. Treatment of axial spondyloarthritis (ankylosing spondylitis and nonradiographic axial spondyloarthritis) in adults. UpToDate [database online]. Waltham, MA. Last updated August 24, 2022. Available at: <http://www.uptodate.com/home/index.html>.
27. UpToDate, Inc. Treatment of peripheral spondyloarthritis. UpToDate [database online]. Waltham, MA. Last updated March 17, 2022. Available at: <http://www.uptodate.com/home/index.html>.
28. Menter A, Gelfand JM, Connor C, et al. Joint American Academy of Dermatology–National Psoriasis Foundation guidelines of care for the management of psoriasis with systemic nonbiologic therapies. *Journal of the American Academy of Dermatology*. 2020;82(6):1445-1486.
29. Menter A, Strober BE, Kaplan DH, et al. Joint AAD-NPF guidelines of care for the management and treatment of psoriasis with biologics. *Journal of the American Academy of Dermatology*. 2019;80(4):1029-1072.

30. Rubin DT, Ananthakrishnan AN, Siegel CA, Sauer BG, Long MD. ACG Clinical Guideline: Ulcerative Colitis in Adults. *Am J Gastroenterol*. 2019;114(3):384-413.
31. Feuerstein JD, Isaacs KL, Schneider Y, et al. AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology*. 2020;158(5):1450-1461. doi:10.1053/j.gastro.2020.01.006
32. Paschos P, Katsoula A, Salanti G, et al. Systematic review with network meta-analysis: the impact of medical interventions for moderate-to-severe ulcerative colitis on health-related quality of life. *Aliment Pharmacol Ther*. 2018 Dec;48(11-12):1174-1185. doi: 10.1111/apt.15005. Epub 2018 Oct 30. PMID: 30378141.
33. Trigo-Vicente C, Gimeno-Ballester V, García-López S, et al. Systematic review and network meta-analysis of treatment for moderate-to-severe ulcerative colitis. *Int J Clin Pharm*. 2018 Dec;40(6):1411-1419. doi: 10.1007/s11096-018-0743-4. Epub 2018 Nov 26. PMID: 30478492.
34. Bonovas S, Lytras T, Nikolopoulos G, et al. Systematic review with network meta-analysis: comparative assessment of tofacitinib and biological therapies for moderate-to-severe ulcerative colitis. *Aliment Pharmacol Ther*. 2018 Feb;47(4):454-465. doi: 10.1111/apt.14449. Epub 2017 Dec 4. PMID: 29205421.
35. Turner et al. Management of Paediatric Ulcerative Colitis, Part 1: Ambulatory Care—An Evidence-based Guideline From European Crohn's and Colitis Organization and European Society of Paediatric Gastroenterology, Hepatology and Nutrition, *Journal of Pediatric Gastroenterology and Nutrition*: August 2018.
36. American Academy of Dermatology – Guidelines of Care for the Management of Atopic Dermatitis (2014). Available at: <https://www.aad.org/member/clinical-quality/guidelines/atopic-dermatitis>
37. Eichenfield L, Tom, W., Chamlin S., et al. Guidelines of care for the management of atopic dermatitis. *JAAD*. 2014 Feb 1. 70(2): 338-351.
38. Singh S, Fumery M, Sandborn WJ, et al. Systematic review and network meta-analysis: first- and second-line biologic therapies for moderate-severe Crohn's disease. *Aliment Pharmacol Ther*. 2018;48(4):394-409. doi:10.1111/apt.14852
39. Ma C, Lee JK, Mitra AR, et al. Systematic review with meta-analysis: efficacy and safety of oral Janus kinase inhibitors for inflammatory bowel disease. *Aliment Pharmacol Ther*. 2019;50(1):5-23. doi:10.1111/apt.15297
40. Nelson SM, Nguyen TM, McDonald JW, et al. Natalizumab for induction of remission in Crohn's disease. *Cochrane Database Syst Rev*. 2018;8(8):CD006097. Published 2018 Aug 1. doi:10.1002/14651858.CD006097.pub3
41. Lichtenstein GR, Loftus EV, Isaacs KL, Regueiro MD, Gerson LB, Sands BE. ACG Clinical Guideline: Management of Crohn's Disease in Adults. *Am J Gastroenterol*. 2018;113(4):481-517.
42. Lamb, Christopher Andrew et al. "British Society of Gastroenterology consensus guidelines on the management of inflammatory bowel disease in adults." *Gut* vol. 68,Suppl 3 (2019): s1-s106. doi:10.1136/gutjnl-2019-318484
43. UpToDate, Inc. Overview of the management of Crohn disease in children and adolescents. UpToDate [database online]. Waltham, MA. Last updated September 14, 2021. Available at: <http://www.uptodate.com/home/index.html>.
44. van Rheenen PF, Aloï M, Assa A, et al. The Medical Management of Paediatric Crohn's Disease: an ECCO-ESPGHAN Guideline Update [published online ahead of print, 2020 Oct 7]. *J Crohns Colitis*. 2020;jjaa161. doi:10.1093/ecco-jcc/jjaa161
45. Feuerstein JD, Ho EY, Shmidt E, et al. AGA Clinical Practice Guidelines on the Medical Management of Moderate to Severe Luminal and Perianal Fistulizing Crohn's Disease. *Gastroenterology*. 2021;160(7):2496-2508. doi:10.1053/j.gastro.2021.04.022
46. King B, Ohyama M, Kwon O, et al. BRAVE-AA Investigators. Two Phase 3 Trials of Baricitinib for Alopecia Areata. *N Engl J Med*. 2022 May 5;386(18):1687-1699.

47. Meah N, Wall D, York K, et al. The Alopecia Areata Consensus of Experts (Ace) study: Results of an international expert opinion on treatments for alopecia areata. *J Am Acad Dermatol.* 2020;83(1):123-130.
48. Seneschal J, Speeckaert R, Taïeb A, et al. Worldwide expert recommendations for the diagnosis and management of vitiligo: Position statement from the international Vitiligo Task Force-Part 2: Specific treatment recommendations. *J Eur Acad Dermatol Venereol.* 2023;37(11):2185-2195.

History

Approved Date	Effective Date	Version	Action and Summary of Changes
04.16.2025	TBD	66.27.00.AH-5	-Added criteria for Opzelura in atopic dermatitis and vitiligo
08.14.2024	03.01.2025	66.27.00.AH-4	Approved by DUR Board - Split 66.27.00 policy into different policies -Added new drug indications when applicable -Update language in medical necessity section
Previous policy changes (relevant from Cytokine & CAM Antagonists Policy)			
Date	Action and Summary of Changes		
10.21.2021	Removed Hyrimoz from the policy and updated the initial dosing for infliximab.		
11.30.2020	Removed Preferred/Non-Preferred listing and added link to AHPDL publication		
11.12.2020	Added language in clinical policy section for cases which do not meet policy criteria		
09.01.2020	Updated wording in clinical criteria for products with only one preferred option.		
08.19.2020	Approved by DUR Board		
8.20.2020	Update to dosing and limits section for all products and indications		
08.12.2020	Updated policy clinical criteria and dosing & quantity limits to include nonradiographic axial spondyloarthritis		
06.01.2020	Added new agents to class; updated age limit for Uveitis indication; updated dosing and quantity limits; updated HCPCS coding		
07.31.2019	Updated criteria that trial of preferred biologics only applies to non-preferred biologics		
06.07.2019	Updates to TB skin test requirements for apremalast; updates to initial authorization clinical criteria		
11.02.2018	Addition of Hyrimoz (adalimumab-adaz)		
09.07.2018	Addition of new medication		
08.16.2017	New Policy		

Cytokine and CAM Antagonists: Janus Associated Kinase (JAK) Inhibitors

Please provide the information below, please print your answer, attach supporting documentation, sign, date, and return to our office as soon as possible. **Without this information, we may deny the request in seven (7) working days.**

Apple Health Preferred Drug list: <https://www.hca.wa.gov/assets/billers-and-providers/apple-health-preferred-drug-list.xlsx>
For policy criteria, see: <https://www.hca.wa.gov/billers-providers-partners/program-information-providers/apple-health-medicaid-drug-coverage-criteria>

Date of request:	Reference #:	MAS:	
Patient	Date of birth	ProviderOne ID	
Pharmacy name	Pharmacy NPI	Telephone number	Fax number
Prescriber	Prescriber NPI	Telephone number	Fax number
Medication and strength		Directions for use	Qty/Days supply

- Is this request for a continuation of therapy? Yes No
- Is this prescribed by, or in consultation with, any of the following? Check all that apply:

<input type="checkbox"/> Allergist	<input type="checkbox"/> Dermatologist	<input type="checkbox"/> Gastroenterologist
<input type="checkbox"/> Immunologist	<input type="checkbox"/> Rheumatologist	<input type="checkbox"/> Other. Specify: _____
- Will the requested medication be used in combination with another Cytokine and CAM medication?
 Yes No
- If request is non-preferred, has patient had treatment with one or more preferred Cytokine and CAM medications on the Apple Health Preferred Drug List (AHPDL) that was ineffective, contraindicated or not tolerated?
 Yes. List each medication and duration of trial:

Medication Name: _____	Duration: _____
Medication Name: _____	Duration: _____
Medication Name: _____	Duration: _____

 No. Explain why a preferred product(s) have not been tried: _____
- What is patient current weight: _____ kg Date taken: _____
- Indicate patient's diagnosis and answer the associated questions as indicated:
 - Alopecia areata (questions 7 - 10)
 - Ankylosing Spondylitis (questions 11 - 15)
 - Atopic dermatitis (questions 16 - 20)
 - Crohn's Disease (questions 21 - 23)
 - Non-radiographic axial spondyloarthritis (questions 11 - 15)
 - Plaque Psoriasis (questions 24 - 28)
 - Polyarticular Juvenile Idiopathic Arthritis (questions 29 - 30)
 - Psoriatic Arthritis (PsA) (questions 31 - 34)

- Rheumatoid Arthritis (questions 35 -37)
- Ulcerative Colitis (questions 38 – 40)
- Vitiligo (questions 41-44)

For diagnosis of Alopecia areata

7. Is patient's current episode of alopecia areata lasting more than 6 months? Yes No
8. Has patient had $\geq 50\%$ of the scalp hair loss (Severity of Alopecia Tool [SALT] score $>50\%$)? Yes No
9. Does patient have a history of failure, contraindication, or intolerance to any of the following? Check all that apply:
- High-potency topical corticosteroids [minimum trial of 6 weeks]
 - Intralesional corticosteroids [minimum trial of 6 weeks]
 - Systemic therapy (i.e., oral corticosteroids, methotrexate, cyclosporine) [minimum trial of 6 weeks].
10. **For continuation of therapy:** Has documentation been submitted demonstrating a positive clinical response (i.e. improvement in SALT score)?
- Yes No

For diagnosis of Ankylosing Spondylitis or Non-radiographic axial spondyloarthritis:

11. Does patient have high disease activity as indicated by one of the following?
- Bath Ankylosing Disease Activity Index (BASDAI) score of at least 4
 - Ankylosing Spondylitis Disease Activity Score (ASDAS) score of at least 2.1
12. Has patient had treatment with at least two different NSAIDs that has been ineffective, contraindicated or not tolerated [minimum trial of four weeks]? Yes No
13. Has patient's disease manifested as one of the following?
- Axial disease Peripheral arthritis
14. Has patient had treatment with at least one non-Cytokine and CAM disease-modifying antirheumatic drug (DMARD) that has been ineffective, contraindicated or not tolerated [minimum trial of 3 months]?
- Yes No
15. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response (e.g., decrease in BASDAI or ASDAS score)? Yes No

For diagnosis of Atopic dermatitis:

16. Indicate disease severity for patient. Check all that apply:
- Body surface area (BSA) involvement of at least 10%
 - Body surface area (BSA) involvement does not exceed 20%
 - Involvement of sensitive skin areas such as hands, feet, face, neck, genitalia, or intertriginous areas
 - Baseline disease severity scale scoring supporting diagnosis of moderate to severe chronic atopic dermatitis (e.g., Investigator's Global Assessment (IGA) score of 3 or greater; Eczema Area and Severity Index (EASI), Patient Oriented Eczema Measure (POEM), etc.)
 - Other. Explain:
17. Indicate if patient is experiencing functional impairment, due to atopic dermatitis, of any of the following. Check all that apply:

- Activities of daily living (ADLs)
- Skin infections
- Sleep disturbances
- Other. Specify:

18. Has patient had a history of failure, defined as the inability to achieve or maintain remission, to any of the following, unless all are contraindicated or clinically inappropriate [minimum trial of 28-days each]? Check all that apply:
- Topical corticosteroids of at least medium/moderate potency (e.g. clobetasol, betamethasone, halobetasol)
 - Topical calcineurin inhibitors (e.g. pimecrolimus cream, tacrolimus ointment)
 - Topical PDE-4 inhibitors (e.g. crisaborole)
 - All are contraindicated or clinically inappropriate. Explain:
19. **For abrocitinib (Cibinqo) and upacitinib (Rinvoq)**, has treatment with dupilumab (Dupixent) has been ineffective, contraindicated, or not tolerated [minimum trial of 16 weeks]? Yes No
20. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response as defined by any of the following? Check all that apply:
- Reduction in body surface area involvement of at least 20% from baseline
 - Achieved or maintained clear or minimal disease from baseline (equivalent to IGA score of 0 or 1)
 - Experienced or maintained a decrease in EASI score of at least 50% from baseline
 - Improvement in functional impairment (e.g., improvement in ADLs, skin infections, or sleep disturbance)

For diagnosis of Crohn's Disease (CD)

21. Has treatment with any of the following conventional therapies that have been ineffective, contraindicated, or not tolerated? Check all that apply:
- Oral corticosteroids (e.g., prednisone, methylprednisolone) used short-term to induce remission or alleviate signs/symptoms of disease flare
 - Immunomodulatory agent (e.g., methotrexate, azathioprine, 6-mercaptopurine) [minimum trial of 12 weeks]
22. Does patient have documentation of high-risk disease (e.g., symptoms despite conventional therapy, obstruction, abscess, stricture, phlegmon, fistulas, resection, extensive bowel involvement, early age of onset, growth retardation, Crohn's Disease Activity Index (CDAI) > 450, Harvey-Bradshaw index > 7)? Yes No
23. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response (e.g., improvement in endoscopic activity, taper or discontinuation of corticosteroids, reduction in number of liquid stools, decrease in presence and severity of abdominal pain, decrease in CDAI, decrease in Harvey-Bradshaw index)? Yes No

For diagnosis of Plaque Psoriasis

24. Does patient have presence of ongoing disease for greater than 6 months? Yes No
25. Please indicate the following for patient:
- Disease affects at least 10% body surface area
 - Disease affects the face, ears, hands, feet, or genitalia
26. Have baseline assessments been submitted (e.g., body surface area (BSA), Psoriasis Area and Severity Index (PASI), Psoriasis Physician's Global Assessment (PGA), itch numeric rating scale, etc.)? Yes No
27. Has patient had a history of failure, contraindication, or intolerance to the following? Check all that apply:
- Phototherapy (UVB or PUVA) [minimum trial of 12 weeks]

Treatment with at least one non-Cytokine and CAM DMARD (e.g., methotrexate, cyclosporine, acitretin, azathioprine, etc.) [minimum trial of 12 weeks]

28. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response (e.g., improvement in BSA, PSAI, Psoriasis PGA, itch numeric rating scale)? Yes No

For diagnosis of Polyarticular Juvenile Idiopathic Arthritis

29. Has patient had treatment with at least one non-Cytokine and CAM DMARD (e.g., methotrexate, sulfasalazine, leflunomide, hydroxychloroquine, azathioprine, cyclosporine) that has been ineffective, unless all are contraindicated, or not tolerated [minimum trial of 3 months]? Yes No
30. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response (e.g., improvement in joint pain, swelling, activities of daily living, reduction in diseases flares, etc.)? Yes No

For diagnosis of Psoriatic Arthritis

31. Has patient had treatment with at least one non-Cytokine and CAM disease-modifying antirheumatic drug (DMARD) that has been ineffective, contraindicated or not tolerated [minimum trial of 3 months]? Yes No
32. Does patient have presence of active, severe disease indicated by provider assessment? Yes No
33. Does patient have presence of any of the following? Check all that apply:
- Erosive disease
 - Elevated C-reactive protein (CRP) or erythrocyte sedimentation rate (ESR)
 - Long-term damage interfering with function (e.g., joint deformities, vision loss)
 - Major impairment of quality of life due to high disease activity at many sites (including dactylitis, enthesitis) or functionally limiting arthritis at a few sites.
34. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response (e.g., improvement in joint pain, swelling, activities of daily living, reduction in diseases flares, etc.)? Yes No

For diagnosis of Rheumatoid Arthritis (RA)

35. Have baseline assessments been submitted (e.g., Disease Activity Score for 28 joints (DAS28) with the CRP, DAS28 with ESR, Simplified Disease Activity Index (SDAI), Clinical Disease Activity Index (CDAI), Routine Assessment of Patient Index Data 3 (RAPID3), Patient Activity Scale (PAS) II)? Yes No
36. Has patient had treatment with at least one non-Cytokine and CAM DMARD (e.g., methotrexate, sulfasalazine, hydroxychloroquine, leflunomide, cyclosporine, azathioprine) that has been ineffective, unless all are contraindicated, or not tolerated [minimum trial of 3 months]? Yes No
37. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response (e.g. improvement in DAS28 with CRP/ESR, SDAI, CDAI, RAPID3, PAS II scores)? Yes No

For diagnosis of Ulcerative Colitis

38. Have baseline assessments been submitted (e.g., stool frequency, endoscopy results, presence of rectal bleeding, disease activity scoring tool)? Yes No
39. Has treatment with conventional therapy (e.g., systemic corticosteroids, azathioprine, mesalamine, sulfasalazine) been ineffective, unless all are contraindicated, or not tolerated [minimum trial of 12 weeks]? Yes No
40. **For continuation of therapy:** Has documentation been submitted demonstrating disease stability or a positive clinical response (e.g., decreased stool frequency, decreased rectal bleeding, improvement in endoscopic activity, tapering or discontinuation of corticosteroid therapy, or improvement on a disease activity scoring tool)? Yes No

For diagnosis of Vitiligo:

41. Has patient had chronic vitiligo for more than 3 months? Yes No
42. Is patient's total body surface area (BSA) involvement 10% or less? Yes No
43. Has patient had treatment from any of the following categories that has been ineffective or not tolerated? Check all that apply:
 Phototherapy
 Topical calcineurin inhibitors (tacrolimus)
 Medium to high potency topical corticosteroids (betamethasone, mometasone, clobetasol, fluocinonide)
44. **For continuation of therapy:** Has documentation been submitted demonstrating a positive clinical response? (e.g. improvement in F-VASI and/or T-VASI score from baseline, or reduction in total BSA involvement) Yes No

CHART NOTES ARE REQUIRED WITH THIS REQUEST

Prescriber signature	Prescriber specialty	Date
----------------------	----------------------	------